




Contents lists available at ScienceDirect

Acta Astronautica

journal homepage: www.elsevier.com/locate/actaastro

Invited Paper

Full-genome study of gene expression in lumbar spinal cord of mice after 30-day space flight on Bion-M1 biosatellite



R.R. Islamov^{a,*}, O.A. Gusev^{b,c,1}, A. Tanabe^{c,2}, M. Terada^{d,3}, O.V. Tyapkina^{a,b,e,4},
K.A. Petrov^e, A.A. Rizvanov^{b,5}, I.B. Kozlovskaya^{f,6}, E.E. Nikolskiy^{a,b,e,7},
A.I. Grigorjev^{f,8}

^a Kazan State Medical University, Butlerov St., 49, 420012 Kazan, Russia^b Kazan (Volga Region) Federal University, Kazan Kremlin St., 18, Russia^c Subio Inc., Amami-shi, Japan^d Japan Aerospace Exploration Agency, Tsukuba, Japan^e Kazan Institute of Biochemistry and Biophysics of Russian Acad. Sci., box 30, 420111 Kazan, Russia^f Institute of Biomedical Problems of Russian Acad. Sci., Horoshovskoe shosse St., 76a, 123007 Moscow, Russia

ARTICLE INFO

Article history:

Received 13 October 2015

Accepted 25 January 2016

Available online 15 February 2016

1. Introduction

Zero-gravity is one of the factors that negatively affect a man in space and it is not a surprise as the evolution of all living things proceeded in a one-G environment. The negative effects of zero-gravity set in while in space, but clinically manifest themselves following the cosmonauts' return to Earth, the usual one-G environment. All the systems of the organism, which adapted to the virtually

weight-free environment, become incapable of regular performance in a one-G environment.

The effects of zero-gravity most strongly manifests in the impaired performance of the locomotor system: the strength and endurance of the muscles decrease, the structure and properties of the bones undergo transformation, and the systems responsible for constructing movements malfunction. This pathological condition is called a "Hypogravitational Locomotor Syndrome" (HLS). The history of long space flights shows that the most effective way of preserving the cosmonauts' performance and preparing them for the return to Earth is a regularly performed complex set of physical exercises in conditions imitating a one-G environment during the flight [1]. However, even diligent performance of specifically designed prophylactic sets of exercises fails to completely curb the development of HLS. It is obvious that the success of long inter-planetary flights will greatly depend on achievements in study of the pathogenesis of the HLS on the molecular, cellular and tissue levels.

Animals that have been in space in a biosatellite or held in an environment imitating zero-gravity play an important role in studies of HLS [2–4]. Such studies have shown HLS develops from an impaired afferent impulse from the limbs.

* Corresponding author. Tel./fax: +7 843 292 7654.

E-mail addresses: islamru@yahoo.com (R.R. Islamov),
oleg@cryptobio.com (O.A. Gusev), tabu@subio.jp (A. Tanabe),
terada.masahiro@jikei.ac.jp (M. Terada),
anti-toxin@mail.ru (O.V. Tyapkina), kpetrov2005@mail.ru (K.A. Petrov),
rizvanov@gmail.com (A.A. Rizvanov),
ikozlovs@imbp.ru (I.B. Kozlovskaya), eenik1947@mail.ru (E.E. Nikolskiy),
grigorjev@imbp.ru (A.I. Grigorjev).

¹ Tel.: +7 190 596 9811; fax: +7 843 292 7347.² Tel.: +81 997 69 3440; fax: +81 997 69 3440.³ Tel.: +81 29 868 3956; fax: +81 29 868 5000.⁴ Tel.: +7 843 292 7647; fax: +7 843 292 7347.⁵ Tel.: +7 843 233 71 82; fax: +7 843 233 9448.⁶ Tel. +7 499 195 2375; fax: +7 499 195 2253.⁷ Tel./fax: +7 843 292 7347.⁸ Tel. +7 499 195 2363; fax: +7 499 195 2253.